Extrinsic feedback for motor learning after stroke: what is the evidence?

Disability and Rehabilitation, Volume 28, No. 13-14, Pages 831-840. doi:10.1080/09638280500534937

Paulette M. van Vliet, PhD Hwwtg'Fellow School of Health Sciences Hcewn{ ''qh'J gcnj Wpkxgtukv{ ''qh'P gy ecuvg.'P UY .'Cwuvtcrkc'' Email: paulette.vanvliet@pgy ecuvgQgf wQcw'

Please address correspondence to Dr. van Vliet

Gabriele Wulf, PhD

Department of Kinesiology

University of Nevada, Las Vegas

e-mail: gabriele.wulf@ccmail.nevada.edu

...

- ...
- ...

Abstract

Extrinsic feedback for motor learning after stroke: what is the evidence?

Purpose There is little guidance on using extrinsic feedback to enhance motor learning after stroke. This narrative review synthesises research findings and identifies questions remaining to be answered.

Method A summary is given relating to the use of extrinsic feedback in healthy subjects. Then, research concerning content of feedback, feedback scheduling, and attentional focus is discussed in relation to patients with stroke.

Results Though research is scarce, preliminary key findings were as follows: Patients' balance performance can improve from receiving visual feedback about weight distribution during practice; auditory feedback of force production may improve performance of sit-to-stand; providing feedback on less than 100% of trials, and giving summary or average feedback may enhance learning; instructions or feedback inducing an external focus may be more effective than those with an internal focus. Further research is needed concerning the relative benefits of verbal, visual, video and kinematic feedback; reduced feedback frequencies and summary feedback schedules; feedback delays, error estimation, and self-controlled feedback; and attentional focus of feedback.

Conclusions Although there are some indications that feedback might enhance motor learning after stroke, there are many areas as yet not examined and there is clearly a need for considerable research in this area.

Extrinsic feedback for motor learning after stroke: what is the evidence?

1. Introduction

Feedback can enhance motor learning in healthy subjects. During training, the performer uses feedback to detect errors in performance by comparison of their movement to the expected goal, in order to improve the next attempt. Careful planning of content, scheduling and attentional focus induced by the feedback can enhance the effectiveness of training considerably. However, research examining these issues in stroke patients is scarce. This narrative review aims to synthesise the findings from current research on delivery of feedback to people with stroke, and identify the main questions that remain to be answered. Firstly, a short introduction will be given placing feedback in the context of motor learning and summarising findings relating to the content and scheduling of feedback, as well as effects of attentional focus of feedback in the learning of healthy subjects.

Feedback can be classified as either 'intrinsic' or 'extrinsic'. Intrinsic feedback refers to a person's own sensory-perceptual information that is available as a result of movement being performed. Several sensory processes, including vision, proprioception, touch, pressure and audition, can mediate this information. Intrinsic feedback helps to formulate a person's internal representation of the movement goal he or she is trying to achieve. Extrinsic (or 'augmented') feedback, is feedback given in addition to intrinsic feedback, usually from an outside source, such as when a therapist comments on the quality of a stroke patient's attempt to stand from a chair. Extrinsic feedback has been categorised into either 'knowledge of results' (KR) or 'knowledge of performance' (KP). KR is 'externally presented information about the outcome of performing a skill or about achieving the goal of the performance' [1]. For example, a patient might be told that

she was able to transfer 40% of body weight on to the hemiplegic leg, with bathroom scales being used to measure the result. KP is 'information about the movement characteristics that led to the performance outcome' [1]. For example, the therapist might tell the patient that the knee needs to be more extended in order to bear more weight on the hemiplegic leg.

Although it is clearly possible to learn motor tasks without extrinsic feedback, comparisons of KR and KP to no feedback at all show improved retention with feedback [2,3]. Feedback that provides information on errors in movement and how to correct them, can facilitate achievement of the movement goal by increasing the level of skill attained or by speeding up the learning process [1]. In the case of KP, this information may be more difficult to obtain intrinsically, as learners may not be so aware or so knowledgeable about their movements as the trainer, or may not be able to see the movements while actually performing the task.

Feedback is also important for sustaining motivation during learning. Extrinsic feedback can encourage persistence to master a skill in a physical education situation [4]. Skill acquisition is generally assumed to be facilitated more effectively if feedback contains information about errors rather than correct performance [1] (although see Chiviacowsky and

Wulf [5]). Knowledge of correct performance can be helpful for motivation.

2. Feedback for learning in healthy subjects

As a preamble to the review of evidence on the use of feedback with people with stroke, the available information on content, scheduling and attentional focus of feedback with healthy subjects is summarised here.

Content

Prescriptive feedback (describing the errors *and* suggesting how to correct them [6]) was found to be more effective than descriptive feedback (just describing the errors) [7]. Increased precision of KR feedback can lead to better learning, though the research is inconsistent [8]. Thorpe [3] argues that to determine the content of feedback, the trainer needs to have analysed the movement task and identified all the essential components that produce skilled performance of the task. Feedback can then be given to the learner on these components. Feedback is commonly given verbally by the trainer, or visually (e.g. demonstration, video), or by manual assistance.

Verbal feedback can be an effective tool and erroneous verbal feedback can even override the person's own correct visual feedback in some situations. In a study by Beukers et al [9], subjects practised an anticipation timing task. When given erroneous verbal feedback, it had the effect of overriding the person's own visual feedback, such that subjects adjusted their response to the incorrect verbal feedback. Also, verbal KR may be redundant when KR is inherent in a task. The same study demonstrated that there was no change in performance if correct verbal KR was given, compared to just receiving the visual feedback with each attempt.

Visual feedback is commonly given by demonstration by the trainer or by videotape. For more experienced learners, self evaluation of videotaped performance works well as shown in a study of tennis players [2]. For less experienced learners, attention-directing cueing can be helpful, however [7,10]. A review of videotape studies by Rothstein and Arnold [10] concludes that videotape should be used over a number of weeks to enhance its effectiveness as a learning tool. Kinematic feedback has been found to be effective for learning of a bimanual coordination task [11], learning a golf shot [12] and increasing power output during a leg press exercise [13]. It frequently involves displaying results graphically to the learner, for example as joint angular displacement. Biofeedback provides information about physiological processes through the use of instrumentation [1].

Scheduling

Feedback can have negative effects if provided too frequently, such as the learner becoming dependent on it, or movement instability [8,14]. Studies have assessed various reduced feedback frequency manipulations, bandwidth, summary or average feedback manipulations, delayed feedback and error estimation procedures, as well as self-controlled feedback schedules.

Regarding reduced feedback frequency, a number of studies have shown that reducing the proportion of trials (e.g. 50 %) for which feedback is presented during the practice phase can result in more effective learning than presenting feedback after every trial (100%) (e.g. Lai et al [15], Weeks et al [16], Winstein et al [17], Wulf et al, [18], Wulf et al [1 9]. That is, when performance is compared on retention or transfer tests (without or with feedback), groups that received less feedback during practice typically show enhanced learning. Learning benefits resulting from a reduced feedback frequency are may also be due to increased movement stability during practice [15,20].

In both summary feedback and average feedback conditions, performers are provided with feedback about a set of trials (e.g. 5) after this set has been completed. Whereas summary feedback involves feedback about every trial in the set, average feedback refers to the average performance on that set of trials. Beneficial effects of summary feedback relative to single-trial feedback have been found in a number of studies [21-23]. The optimal number of t rials summarized depends on the complexity of the task in relation to the performer's skill level, with shorter summary lengths being more effective for more difficult tasks or less experienced performers [21, 23, 24]. Results for average feedback are more equivocal [23,25,26].

Bandwidth feedback involves qualitative feedback when performance is within a certain range of error (e.g., 10%), indicating that performance is "correct", and quantitative feedback when performance is outside the bandwidth (e.g., 70 ms too fast). Several studies have demonstrated benefits of bandwidth feedback relative to (quantitative) feedback provided after each trial [20, 27, 28]. The primary advantage of bandwidth feedback for learning seems to be

that it promotes movement stability during practice. As a consequence, bandwidth feedback often results in more stable retention performance as well.

A feedback manipulation that has consistently been found to be detrimental to learning is feedback that is concurrently provided with the movement. Even though concurrent feedback has very strong performance-enhancing effects when it is present during practice, it typically results in clear performance decrements when it is withdrawn in retention or transfer tests, re lative to feedback presented after the movement [29-32]. Also, presenting feedback instantaneously after the completion of the movement tends to create a strong dependency on it. One reason for the detrimental effects of concurrent and instantaneous feedback might be that they prevent spontaneous errors estimations, based on the processing of intrinsic feedback, that might occur during or after the movement. In contrast, delaying the feedback for a few seconds seems to promote those error estimations and has indeed been found to produce more effective learning [33]. Along the same lines, studies in which participants were specifically instructed to estimate their errors also provide evidence for the learning benefits of subjective movement evaluations [33,34].

Allowing the performer to decide when he or she wants to be provided feedback can be beneficial for learning [36-38]. It has been suggested that the perception of self-control enhances learning because it leads to a more active involvement of the learner in the learning process, promotes a deeper information processing [39-41], is more motivating [42,43] and makes performers take charge of their own learning process (e.g. Ferrari [44]).

Focus of attention

Numerous studies have shown that instructions that direct individuals' attention to the effects of their movements on the environment (e.g., implement, apparatus), thereby inducing a so-called "external" focus of attention, are more effective for learning than instructions that direct attention to the movements themselves, or inducing an "internal" focus (for a review, see Wulf & Prinz, [45]). In a study by Shea and Wulf [46], a balance task (stabilometer) was used that required

participants to keep the platform on which they were standing horizontal. Participants were presented with concurrent visual feedback, which consisted of the platform movements displayed on a computer screen. While one group was informed that the feedback represented movements of their feet (internal focus), the other group was told that the feedback represented lines attached to the platform in front of each of the performer's feet (external focus). The results showed that the group that interpreted the feedback as "external" demonstrated more effective balance than the group that interpreted it as "internal". Interestingly, the beneficial effects of external focus feedback were not only seen during practice when the feedback was present, but also in retention when the feedback was withdrawn.

The benefits of adopting an external focus of attention have been explained with a more automatic type of movement control that is promoted when attention is directed to the movement effect. In contrast, when participants are asked to focus on their movements, they tend to actively intervene in the motor control processes, thereby disrupting automatic control processes ("constrained action hypothesis"). Support for this notion comes from studies showing shorter probe reaction times (indicating reduced attentional demands and a greater degree of automaticity), faster and more reflexive movement adjustments, and reduced electromyographic (EMG) activity for individuals adopting an external compared to an internal focus [47-49].

In contrast to the somewhat "artificial" situation in the Shea and Wulf [46] study, physiotherapists typically provide patients with verbal feedback that refers to the aspect of the movement that needs the most improvement. This type of feedback was examined in a study by Wulf, McConnel, Gärtner, and Schwarz [50]. Specifically, they examined the effectiveness of feedback for the learning of motor skills (i.e., volleyball "tennis" serve, lofted soccer kick), with feedback either being worded in a way that it induced an internal focus (e.g., "shift your weight from your back leg to your front leg") or external focus (e.g., "shift your weight toward the target"). The results showed that learning was enhanced by the external relative to the internal focus feedback. Regarding frequency of feedback, in this study it was found that 100% external

focus feedback was no different from 33% external focus feedback, and even tended to be more effective, in contrast to 100% internal focus feedback, which was detrimental to learning relative to 33% feedback [50]. This suggests that frequent feedback is not detrimental (and perhaps even beneficial) if it induces an external focus. Wulf et al. [50]. argued that a high frequency of internal-focus feedback might be detrimental because it encourages learners to focus *too much* on their movements, while these effects should be attenuated under reduced feedback conditions. In contrast, feedback inducing an external focus presumably promotes a more automatic type of control, and frequent reminders do not appear to hinder the learning process.

3. Feedback for people after stroke

An important question for physiotherapists working in stroke rehabilitation is whether the research findings for healthy subjects apply to the person with stroke. This depends partly on whether people with stroke learn in the same way as those with intact nervous systems, and whether the tasks studied in the research are similar to those which the patient needs to learn. There is a small amount of research to answer the first consideration. After stroke, intrinsic feedback systems may be compromised, making it difficult for the person to determine what needs to be done to improve performance. Extrinsic feedback may thus be even more important to people with stroke. Studies have shown that patients with unilateral stroke are able to learn new motor skills [51,52], however implicit motor learning (learning perceptual-motor skills by physical practice without conscious awareness) may be impaired [53] particularly in patients with temporal lobe damage [54]. It has been suggested that provision of KR may allow explicit memory (knowledge of facts, events, and episodes) to assist motor learning in these patients [53,55]. Secondly, the skills that need to be learned by people with stroke include the performance of everyday tasks such as standing up, reaching to grasp an object, and walking. These seem similar to the kinds of sport skills that have been the focus of much of the research in healthy subjects. Although standing up may not seem as difficult as a sports skill such as serving

in tennis, it may be perceived as difficult by the person who has suffered a stroke. Parts of tasks, e.g. extension of the wrist, are also practised, which also resemble some of tasks studied in healthy subjects (e.g. Beukers [9]). Although the processes by which people with stroke learn need to be further elucidated with more research, there is some indications that findings from healthy subjects are relevant for this group.

Feedback content should be adjusted for the stage of learning of the subject. In the early stages of learning, information about the general framework and sequencing of components is beneficial, whereas in a later stage feedback can be more precise [56]. The role of the therapist is to provide feedback that is likely to assist learning in the most effective way. Feedback is probably delivered intuitively much of the time during stroke rehabilitation [57]. However, there is some evidence to guide the therapist so that feedback can be used in a more focused and deliberate way. In the next section, the evidence to date on use of extrinsic feedback in people with stroke is presented and critically appraised.

Review of evidence of use of feedback for learning after stroke

Content of feedback

Existing research on content of feedback for stroke patients focuses on assessing the effectiveness of instrumental feedback devices, rather than more fundamental issues such as contrasts between KR and KP, descriptive and prescriptive feedback, and different content of verbal feedback for different stages in the learning process. One exploratory study which included subjects with stroke, investigated the socio-affective characteristics of extrinsic feedback in physiotherapy [57]. Patient-physiotherapist interactions were videotaped and then subjected to systematic observation, in which behaviour was recorded into pre-determined categories of verbal and physical communications. This study found that verbal feedback was used extensively by physiotherapists and was used more frequently than visual feedback [57]. The verbal feedback was mainly motivational and reinforcing, with information feedback being used rarely. It was sometimes given concurrently with the patient's practice of the movement, which research in healthy subjects has indicated is counterproductive to learning [29-32]. This study does not allow assessment of the effect of different feedback on learning since it is exploratory – this should be a target of further research.

One study has specifically investigated the effect of KR after stroke [58]. Stroke subjects in this study were randomised into 3 groups and then underwent a 3-week training programme of upper limb tasks. The first group received the training with KR, the second received the training without KR and the third group did not have the training. Although the training itself produced significant results compared to no training, when performance was measured at the end of the 3 weeks, there was no substantial extra effect for KR. This could be because the tasks chosen had inherent information about the movement outcome, e.g. hitting targets with a stylus and placing objects on top of other objects, so that additional KR, which took the form of bar diagrams on a computer screen, did not enhance learning further. This is a similar result to the study by Beukers et al [9] with healthy subjects, who found that extra verbal KR was redundant when outcome information is inherent in a task. Therefore, when outcome information is available, we can make the cautious interpretation that therapists may not need to give further informational KR. It may have a motivational role, however, and this needs to be investigated. Studies are also needed to elucidate whether KR will be useful in tasks where KR is not inherent, with stroke subjects. Therapists should also consider that setting tasks with inherent KR could be desirable, as it encourages selfcontrol of performance, which has been found to enhance learning in healthy subjects [39-41]. Verbal feedback has been an element of treatment for stroke in two other randomised controlled trials, where positive effects were found for the entire intervention [59,60]. However, it was not possible to discern the particular contribution of the verbal feedback in these multi-faceted treatments.

11

Several studies have demonstrated that stroke patients can benefit from receiving visual feedback about weight distribution and weight shift activity. In a randomised controlled trial, Sackley and Lincoln [61] found significant improvements compared to a control group in stance symmetry and sway, motor function and activities of daily living after a 4-week training programme in which patients received feedback on a computer screen while attempting balance tasks on a balance platform. These changes were present 4 weeks after the intervention, but not at 12 weeks. In a similar study in which visual feedback was given about centre of gravity to more acute stroke patients [62], balance performance also improved but was not significantly different to a group receiving the same amount of balance training via the verbal and tactile cues more typically used in physiotherapy. Altschuler et al [63] compared training of symmetric arm movements in a mirror to symmetric arm movements using a plastic sheet in a randomised crossover design. Subjective comments by the patients and ratings of improvement by researchers based on videotapes of 'cardinal movements of the upper limb' indicated that it might be beneficial for some patients. The lack of reliable outcome measures and adequate statistical analysis, however, means that this finding should be regarded with caution. Visual feedback was concurrently and continuously displayed to the learner in the above studies, which could may be detrimental to learning, even though it improves performance, because of reduced opportunity for the patient to use their own intrinsic feedback, leading to a potential dependency on extrinsic feedback [64]. In future studies comparing different content of feedback, it would be useful if the scheduling of feedback reflected the recommendations from more recent research, so that it was slightly delayed and given with reduced frequency.

Auditory feedback of force production was also shown to improve performance of the sit-to-stand movement by Enghardt et al [65]. In a randomised controlled trial, the experimental group received ground reaction force feedback via auditory input, whereas the control group received no feedback about ground reaction force. When measured at the end of a 5-week programme of daily practice of sit-to-stand, the experimental group achieved significantly greater symmetry in bodyweight distribution than the control group. Portable weight distribution auditory feedback devices have been employed also (e.g. Batavia et al [66]). One study evaluated the effect of such a device on performance of sit-to-stand [67]. A small group of patients receiving auditory feedback during practice of sit-to-stand for a period of 3 weeks were compared to a control group who underwent the practice without the feedback. The results were equivocal although there was a tendency for the feedback group to increase force production through the affected lower limb more than control subjects. Dursun et al [68] also demonstrated positive effects for an angular biofeedback device for sitting balance. The device was positioned on the midline of the back, and gave information about tilting from the erect position with feedback given both by visual and auditory signals. Balance performance was significantly better on some of the measures used, compared to the control group, after 10 days of treatment, but the difference was not maintained at discharge. The study was not randomised and so it is not clear whether biased patient selection contributed to the results.

Effects of kinematic feedback in stroke rehabilitation have received little attention in the literature. One quasi-randomised study has examined the effect of kinematic feedback via electrogoniometry for the purpose of limiting knee hyperextension [69]. Peak knee hyperextension was improved more than the control group after a 4-week training period. A positive aspect of this study is that frequency of feedback information from research in healthy subjects was applied so that patients only received feedback if the knee was extended past the 0 degree position (bandwidth feedback). It has been argued that another group receiving non-bandwidth feedback could have been included to assess the effect of bandwidth [64]. This is a good point as it is not clear yet whether the general finding that reduced frequency of feedback is better for retention in healthy subjects, applies to subjects with stroke. It is particularly relevant, as the electrogoniometry feedback induces an internal focus of attention (see section on healthy

subjects). Another admirable aspect of the design was that results were based on a retention test 4 weeks after cessation of the training, unlike some of the studies with people with stroke, which tested the change in performance as a result of the feedback intervention. An interesting case study showed improvement in gait parameters when regularly shown a graph of performance [70]. The patient was covertly monitored by staff who observed the size of her base of support and step length, and this information was fed back to the patient daily on a graph showing the number of times the correct step length or base of support was observed. Another common form of biofeedback, EMG, will not be discussed here, as it has been the subject of several earlier papers.

There are other instrumental forms of feedback under development for stroke rehabilitation, such as virtual reality-augmented training. For example, the patient can wear different types of gloves containing infrared sensors and force transducers while attempting upper limb movements – one for monitoring amplitude, speed and fractionation of movement, the other, a force feedback glove, monitoring strength of finger flexion and extension movements [71,72]. Online visual, auditory and force feedback was provided via a personal computer. Exercises were in the form of imaginative computer games with graphics feedback and goals were set according to ability. One and a half hours of virtual reality training plus practice of fine motor tasks, spread over a period of 3.5 hours per day for 2 weeks, was found to have good effects. A similar training programme which used video-game like tasks had good effects for training pronation, supination, and wrist movements [73]. A type of web-based telerehabilitation has been described [74], which aims to provide low cost, intensive, repetitive practice of functional movements via input devices such as a force feedback joystick and traditional mechanical mice, or gyroscopic or force feedback mice. Randomised controlled trials are yet to be conducted to assess the effectiveness of these methods.

Scheduling of feedback

Relatively few studies have examined feedback scheduling effects in stroke patients. Even though no studies seem to have looked at error estimation and self-controlled feedback effects, some studies have examined the effects of feedback frequency, summary, average and summary feedback, as well as feedback delay (for a review, see Ezekiel et al [64]).

Reduced feedback frequency.

Winstein, Merians, and Sullivan [52] used a task, in which participants had to produce a spatially and temporally defined arm movement pattern (lever positioning), to compare feedback frequency effects (100% vs. 67%) in healthy, older adults and a group of individuals who were an average of 2 years post-stroke. In a retention test, the reduced feedback frequency during practice resulted in increased consistency for healthy and stroke patients, even though movement accuracy did not differ from the respective 100% feedback conditions. In a study using a linear arm positioning task [75], different groups of individuals with brain injury and aged-matched controls practiced under 33%, 67%, and 100% feedback conditions. There were no significant differences between feedback frequency groups in retention test, although both reduced frequency groups tended to be more accurate than the 100% condition. Finally, Saladin, Baghdady, and Nichols [76] report that stroke patients practicing an iso metric force modulation task benefited from a reduced feedback frequency (50%) compared to 100% feedback, as demonstrated in a retention test after 24 hours. Thus, even though there is no compelling evidence that reducing the feedback frequency is beneficial for stroke patients, at the very least, the findings suggest that it is not harmful.

Summary and average feedback.

A study by Croce, Horvat, and Roswal [77], using a coincidence timing task, provided some evidence for the effectiveness of summary and average feedback in individuals with traumatic brain injury. Compared to groups that received no feedback (control) or feedback after every trial, both summary and average feedback groups performed more effectively on an immediate retention test, and the summary feedback group was most accurate on a 24-hour retention test.

Overall, the findings regarding feedback scheduling effects in stroke patients are somewhat equivocal. Clearly, more studies are needed, including ones that use more complex and functional tasks (see Ezekiel et al [64]). Also, one factor that needs to b e considered in this context is the relative task difficulty. Wulf and Shea [50,78,79] have argued that, whereas the learning of simple tasks might be enhanced by making practice more difficult or challenging for the learner (e.g., by reducing the feedback frequency), the learning of relatively complex skills might not benefit from, and might even be degraded by increasing the demands imposed on the learner. Clearly, a relatively simple task for a healthy subject could be difficult for a stroke patient. Thus, a reduced feedback frequency might not necessarily have the same advantages for stroke patients as it would have for healthy subjects. Furthermore, given the types of tasks that individuals suffering from a stroke learn in physiotherapy setting, feedback might not be as prescriptive as it often is in many simple tasks practiced in laboratory settings. Real-life tasks often require the coordination of different components to produce skilled performance. To improve performance, the learner has to rely on many sources of intrinsic feedback, and the likelihood of patient becoming dependent on the extrinsic feedback and completely neglecting the processing of intrinsic feedback might be reduced.

Focus of attention

While there do not appear to be studies that specifically examined the effectiveness of external versus internal focus feedback in stroke patients, Fasoli, Trombly, Tickle-Degnen, and Verfaellie [80] investigated the effects of external versus internal focus instructions in patients who had a cerebrovascular accident (CVA). Specifically, they used patients and non-impaired control participants to examine how the type of focus affected performance of daily-life tasks. Fasoli and colleagues found that both groups performed various tasks (e.g., taking an apple off a

shelf and putting it into a basket; moving an empty coffee mug from a table onto a saucer) more effectively if given external rather than internal focus instructions. Movement times were shorter and peak velocities were greater on all tasks, suggesting that these patients as well as control participants pre-planned their movements to a greater extent and used more automatic control processes when they focused externally.

A recent study by Landers, Wulf, Wallmann, and Guadagnoli [81] used a different patient population, namely, persons with Parkinson's disease to examine attentional focus effects. Their participants, who also had a history of falls, performed various balance tasks while their postural sway was measured. Landers and colleagues found no differences between attentional focus conditions (internal, external, control) when the tasks were relatively easy, requiring participants to just stand still on a stable surface with their eyes open or closed, respectively. However, when a more difficult sway-referenced support surface was used (i.e., when the platform, on which the participant was standing, moved forward or b ackward in accordance with their center of pressure), external focus instructions, which directed participants' attention to keeping rectangles under their feet horizontal, resulted in significantly less sway than internal focus instructions (attention directed to keeping the feet horizontal), or no focus instructions.

Even though these studies did not directly examine the effectiveness of feedback as a function of attentional focus in patients with stroke, the results are encouraging in that they demonstrate that the external focus benefits generalize to patient populations, including those who suffer from the consequences of a stroke. Together with the learning advantages of external focus feedback shown for healthy participants, these findings suggest stroke patients would likely also benefit from feedback that is phrased in a way that attention is directed to the movement effect, rather to the movements themselves. This appears to be a fruitful direction for future research. Future studies should also use learning paradigms, including the use of retention or transfer tests without feedback, to examine the permanency of the attentional focus effects, if any. If benefits of practicing a skill under external focus conditions were found in situations in which subjects are

not reminded to adopt, or are even prevented from adopting (e.g., Totsika & Wulf, [82]), a specific attentional focus, this could have important implications for physiotherapy settings.

Key findings for stroke therapy:

Because of the small number of studies available, the following summary comments should be regarded as preliminary:

• Additional verbal KR from the therapist may be redundant when outcome information is inherent in the task.

• Patients' balance performance can improve from receiving visual feedback about weight distribution during practice

• Auditory feedback of force production may improve performance of the sit-to-stand movement

- Providing patients with feedback on less than 100% of the trials may enhance learning
- Giving patients summary or average feedback might benefit the learning process
- Instructions or feedback inducing an external focus may be more effective than those

with an internal focus to improve performance of tasks after stroke.

Questions that need to be answered:

How do physiotherapists deliver feedback in current practice (in detail)?

Do patients with stroke benefit more from prescriptive information on errors than on correct

performance?

Do patients with stroke benefit more from KR or KP?

Is their learning enhanced by video and kinematic feedback?

Do patients with stroke benefit the same from feedback after every trial or a reduced feedback

frequency?

Does a reduced feedback frequency improve retention after stroke?

Is concurrent feedback, feedback immediately after performance, or delayed feedback more effective for retention?

Is feedback with an external focus more beneficial for learning after stroke than feedback with an internal focus?

Do patients with stroke benefit from frequent feedback when feedback induced an external focus? How do feedback type and frequency interact with the stage of learning?

Conclusion

Some of the research presented indicates that feedback can enhance motor learning after stroke. So there is some justification for using it in stroke rehabilitation. However, there are many areas as yet not examined and there is clearly a need for considerable research in this area. There is a dearth of studies on how variables, which have been shown to enhance learning in young, nonimpaired populations, affect the re-learning of motor skills in stroke patients. These include the relative benefits of verbal, visual, video and kinematic feedback, reduced feedback frequencies and summary feedback schedules, as well as feedback delays, error estimation, and selfcontrolled feedback. Another fruitful direction for future research might be a further examination of the effects of feedback inducing an external versus internal focus of attention, and how external focus feedback, if shown to be beneficial, can be implemented in physiotherapy settings. Methodologically, studies examining the effect of feedback on motor learning of patients with stroke have not always distinguished between learning and performance. The main focus has been whether performance is improved during training, or immediately after the training. Yet, as the goal of any intervention is improved performance in the long term (i.e., learning), retention or transfer tests conducted some time after the end of training are necessary.

References

- Magill RA. *Motor Learning and Control: Concepts and applications*. Seventh ed. New York: McGraw-Hill; 2003.
- 2. Herbert E, Landin D, Menickelli J. Videotape feedback: what learners see and how they use it. *Journal of Sport Pedagogy*. 1998;4:12-28.
- Thorpe DE, Valvano J. The effects of knowledge of performance and cognitive strategies on motor skill learning in children with cerebral palsy. *Paediatric Physical Therapy*. 2002;14(1):2-15.
- Solmon MA, Boone J. The impact of student goal orientation in physical education classes. *Research Quarterly for Exercise and Sport*. 1993;64(4):418-424.
- Chiviacowsky S, Wulf G. Feedback after good trials enhances learning. Manuscript submitted for publication. 2005.
- 6. Schmidt RA, Wrisberg CA. *Motor learning and performance: a problem-based learning approach.* 2nd ed. Champaign, IL: Human Kinetics; 2000.
- 7. Kernodle MW, Carlton LG. Information feedback and the learning of multipledegree-of-freedom activities. *Journal of Motor Behaviour*. 1992;24:187-196.
- Salmoni AW, Schmidt RA, Walter CB. Knowledge of results and motor learning: A review and critical reappraisal. *Psychological Bulletin*. 1984;95:355-386.
- Beukers MA, Magill RA, Hall KG. The effect of knowledge of results on skill acquisition when augmented information is redundant. *The Quarterly Journal of Experimental Psychology*. 1992;44A(1):105-117.

- 10. Rothstein AL, Arnold RK. Bridging the gap: Application of research on videotape feedback and bowling. *Motor Skills: Theory into Practice*. 1976;1:36-61.
- Swinnen SP, Walter CB, Lee TD, Serrien DJ. Acquiring bimanual skills: Contrasting forms of information feedback for interlimb decoupling. *Journal of Experimental Psychology: Learning, Memory and Cognition.* 1993;19:1321-1344.
- Wood CA, Gallagher JD, Martino PV, Ross M. Alternate forms of knowledge of results: Interaction of augmented feedback modality on learning. *Journal of Human Movement Studies*. 1992;22:213-230.
- 13. Hopper DM. The influence of visual feedback on power during leg press on elite women field hockey players. *Physical Therapy in Sport*. 2003;4(4):182-186.
- Schmidt RA. Frequent augmented feedback can degrade learning: Evidence and interpretations. In: Requin J, Stelmach GE, eds. *Tutorials in motor neuroscience*. Dordrecht: Kluwer Academic Publishers; 1991:59-75.
- Lai Q, Shea CH. Generalized motor program (GMP) learning: Effects of reduced frequency of knowledge of results and practice variability. *Journal of Motor Behaviour*. 1998;30:51-59.
- Weeks DL, Kordus RN. Relative frequency of knowledge of performance and motor skill learning. *Research Quarterly for Exercise and Sport*. 1998;69:224-230.
- Winstein CJ, Schmidt RA. Reduced frequency of knowledge of results enhances motor skill learning. *Journal of Experimental Psychology: Learning, Memory and Cognition.* 1990;16:677-691.

- Wulf G, Schmidt RA. The learning of generalized motor programs: Reducing the relative frequency of knowledge of results enhances memory. *Journal of Experimental Psychology: Learning, Memory and Cognition.* 1989;15:748-757.
- Wulf G, Schmidt RA, Deubel H. Reduced feedback frequency enhances generalized motor program learning but not parameterization learning. *Journal of Experimental Psychology: Learning, Memory and Cognition*, 1993;19:1134-1150.
- 20. Lai Q, Shea CH. Bandwidth knowledge of results enhances generalized motor program learning. *Research Quarterly for Exercise and Sport*. 1999;70:79-83.
- Schmidt RA, Lange C, Young DE. Optimizing summary knowledge of results for skill learning. *Human Movement Science*. 1990;9:325-348.
- 22. Schmidt RA, Young DE, Swinnen S, Shapiro DE. Summary knowledge of results for skill acquisition: Support for the guidance hypothesis. *Journal of Experimental Psychology: Learning, Memory and Cognition.* 1989;15:352-359.
- Yao W-X, Fischman MG, Wang YT. Motor skill acquistion and retention as a function of average feedback, summary feedback, and performance variability. *Journal of Motor Behaviour*. 1994;26:273-282.
- Guadagnoli MA, Dornier LA, Tandy RD. Optimal length for summary knowledge of results: The influence of task-related experience and complexity. *Research Quarterly for Exercise and Sport.* 1996;67:239-248.
- Young DE, Schmidt RA. Augmented kinematic feedback for motor learning. Journal of Motor Behaviour. 1992;24:261-273.
- Wulf G, Schmidt RA. Average feedback degrades parameter learning. *Journal of Motor Behaviour*. 1996;28:371-381.

- 27. Lee TD, Carnahan H. Bandwidth knowledge of results and motor learning. *The Quarterly Journal of Experimental Psychology*. 1990;42:777-789.
- Sherwood DE. Effects of bandwidth knowledge of results on movement consistency. *Perceptual and Motor Skills*. 1988;66:535-542.
- Park J-H, Shea CH, Wright DL. Reduced frequency concurrent and terminal feedback: A test of the guidance hypothesis. *Journal of Motor Behaviour*. 2000;32:287-296.
- Schmidt RA, Wulf G. Continuous concurrent feedback degrades skill learning: Implications for training and simulation. *Human Factors*. 1997;39:509-525.
- VanderLinden DW, Cauraugh JH, Greene TA. The effect of frequency of kinetic feedback on learning an isometric force production task in nondisabled subjects. *Physical Therapy*. 1993;73:79-87.
- Winstein CJ, Pohl PS, Cardinale C, Green A, Scholtz L, Waters CS. Learning a partial-weight-bearing skill: Effectiveness of two forms of feedback. *Physical Therapy.* 1996;76:985-993.
- 33. Swinnen S, Schmidt RA, Nicholson DE, Shapiro DC. Information feedback for skill acquisition: Instantaneous knowledge of results degrades learning. *Journal of Experimental Psychology: Learning, Memory and Cognition.* 1990;16:706-716.
- Guadagnoli MA, Kohl RM. Knowledge of results for motor learning:
 Relationship between error estimation and knowledge of results frequency.
 Journal of Motor Behaviour. 2001;33:217-224.

23

- 35. Liu J, Wrisberg CA. The effect of knowledge of results delay and the subjective estimation of movement form on the acquisition and retention of motor skill. *Research Quarterly for Exercise and Sport.* 1997;68:145-151.
- 36. Janelle CM, Barba DA, Frehlich SG, Tennant LK, Cauraugh JH. Maximising performance effectiveness through videotape replay and a self-controlled learning environment. *Research Quarterly for Exercise and Sport*. 1997;68:269-279.
- 37. Janelle CM, Kim J, Singer RN. Subject-controlled performance feedback and learning of a closed motor skill. *Perceptual and Motor Skills*. 1995;81:627-634.
- Chiviacowsky S, Wulf G. Self-controlled feedback: Does it enhance learning because performers get feedback when they need it? *Research Quarterly for Exercise and Sport.* 2002;73:408-415.
- McCombs ML. Self-regulated learning and achievement: a phenomenological view. In: Zimmerman BJ, Schunk DH, eds. Self-regulated Learning and Academic Achievement: Theory, Research, and Practice. New York: Springer-Verlag; 1989:51-82.
- 40. Watkins D. Student's perceptions of factors influencing tertiary learning. *Higher Education Research Developments*. 1984;3:33-50.
- Chen D, Singer RN. Self-regulation and cognitive strategies in sport participation. International Journal of Sport Psychology. 1992;23:277-300.
- 42. Bandura A. Perceived self-efficacy in cognitive development and functioning. *Educational Psychologist.* 1993;1:117-148.
- Boekaerts M. Self-regulated learning at the junction of cognition and motivation. *European Psychologist.* 1996;1:100-112.

- 44. Ferrari M. Observing the observers: Self-regulation in the observational learning of motor skills. *Developmental Review*. 1996;16:203-240.
- 45. Wulf G, Prinz W. Directing attention to movement effects enhances learning: A review. *Psychonomic Bulletin & Review*. 2001;8:648-660.
- 46. Shea CH, Wulf G. Enhancing motor learning through external-focus instructions and feedback. *Human Movement Science*. 1999;18:553-571.
- Wulf G, McNevin NH, Shea CH. The automaticity of complex motor skill learning as a function of attentional focus. *Quarterly Journal of Experimental Psychology*. 2001;54:1143-1154.
- 48. McNevin NH, Shea CH, Wulf G. Increasing the distance of an external focus of attention enhances learning. *Psychological Research*. 2003;67:22-29.
- 49. Wulf G, Shea CH, Park J-H. Attention in motor learning: Preferences for and advantages of an external focus. *Research Quarterly for Exercise and Sport*. 2001;72:335-344.
- 50. Wulf G, McConnel N. Enhancing the learning of sport skills through externalfocus feedback. *Journal of Motor Behaviour*. 2002;34(2):171-182.
- Hanlon RE. Motor learning following unilateral stroke. Archives of Physical Medicine and Rehabilitation. 1996;77:811-815.
- Winstein CJ, Merians AS, Sullivan KJ. Motor learning after unilateral brain damage. *Neuropsychologia*. 1999;37:975-987.
- Boyd L, Winstein CJ. Implicit motor-sequence learning in unilateral stroke: Impact of practice and explicit knowledge. *Neuroscience Letters*. 2001;298:65-69.

- 54. Reber PJ, Squire LR. Encapsulation of implicit and explicit memory in sequence learning. *Journal of Cognitive Neuroscience*. 1998;10:248-263.
- 55. Winstein C, Wing AM, Whitall J. Motor control and learning principles for rehabilitation of upper limb movements after brain injury. In: Grafman J, ed. *Handbook of Neuropsychology*. Vol 9. Amsterdam: Elsevier; 2005.
- 56. Gentile AM. Skill acquisition: action, movement and neuromotor processes. In: Carr JH, Shepherd RB, eds. *Movement Science: Foundations for Physical Therapy in Rehabilitation*. London: Heinemann Physiotherapy; 1987.
- 57. Talvitie U. Socio-affective characteristics and properties of extrinsic feedback in physiotherapy. *Physiotherapy Research International*. 2000;5(3):173-188.
- 58. Platz T, Winter T, Muller N, Pinkowski C, Eickhof C, Mauritz K-H. Arm ability training for stroke and traumatic brain injury patients with mild arm paresis: a single-blind, randomized, controlled trial. *Archives of Physical Medicine and Rehabilitation*. 2001;82:961-968.
- Sunderland A, Tinson D, Bradley L, Fletcher D, Hewer RL, Wade DT. Enhanced physical therapy for arm function after stroke. *Journal of Neurology, Neurosurgery and Psychiatry.* 1992;55:530-535.
- Langhammer B, Stanghelle JK. Bobath or Motor Relearning Programme? A comparison of two different approaches of physiotherapy in stroke rehabilitation: a randomized controlled study. *Clinical Rehabilitation*. 2000;14:361-369.
- 61. Sackley CM, Lincoln NB. Single blind randomized controlled trial of visual feedback after stroke: Effects on stance symmetry and function. *Disability and Rehabilitation*. 1997;19:536-546.

- Walker C, Brouwer BJ, Culham EG. Use of visual feedback in retraining balance following acute stroke. *Physical Therapy*. 2000;80(9):886-895.
- 63. Altschuler EL, Wisdom SB, Stone L. Rehabilitation of hemiparesis after stroke with a mirror. *Lancet.* 1999;353:2035-2036.
- 64. Ezekiel HJ, Lehto NK, Marley TL, Wishart LR, Lee TD. Application of motor learning principles: the physiotherapy client as a problem-solver. III Augmented feedback. *Physiotherapy Canada*. 2001;53(1):33-39.
- 65. Engardt M, Ribbe T, Olsson E. Vertical ground reaction force feedback to enhance stroke patients' symmetrical body-weight distribution while rising/sitting down. *Scandinavian Journal of Rehabilitation Medicine*. 1993;25:41-48.
- 66. Batavia M, Gianutsos JG, Kambouris M. An augmented auditory feedback device. *Archives of Physical Medicine and Rehabilitation*. 1997;78:1389-1392.
- 67. Fowler V, Carr J. Auditory feedback: effects on vertical force production during standing up following stroke. *International Journal of Rehabilitation Research*. 1996;19:265-269.
- Dursun E, Hamamci N, Donmez S, Tuzunalp O, Cakci A. Angular biofeedback device for sitting balance of stroke patients. *Stroke*. 1996;27:1354-1357.
- Morris ME, Matyas TA, Bach TM, Goldie PA. Electrogoniometric feedback: Its effect on genu recurvatum in stroke. *Archives of Physical Medicine and Rehabilitation*. 1992;73:1147-1154.
- Canning C, Adams R. Covert monitoring to promote consistency of walking performance: a case study. *Australian Journal of Physiotherapy*. 1985;31(4):152-154.

- Jack D, Boian R, Merians AS, et al. Virtual reality-enhanced stroke rehabilitation.
 IEEE Transactions on Rehabilitation Engineering. 2001;9(3):308-318.
- 72. Merians AS, Jack D, Boian R, et al. Virtual Reality-augmented rehabilitation for patients following stroke. *Physical Therapy*. 2002;82(9):898-915.
- Hermsdorfer J, Marquardt C, Heiss J. Evaluation of a feedback-based training of cerebral hand function deficits using kinematic analysis. *Physikalische Mediziin Rehabilitationmedizin.* 2004;14(4):187-194.
- 74. Reinkensmeyer DJ, Pang CT, Nessler JA, Painter CC. Web-based telerehabilitation for the upper extremity after stroke. *IEEE Transactions on RehabIilitation Engineering*. 2002;10(2):102-108.
- 75. Thomas DM, Harro CC. Effects of relative frequency of knowledge of results on brain injured and control subjects learning a linear positioning task. *Neurology Report.* 1996;20:60-62.
- Saladin LS, Baghdady M, Nichols L. The effects of reduced relative frequency of feedback on motor learning in stroke patients. *Physical Therapy*. 1994;5:S122.
- 77. Croce R, Horvat M, Roswal G. Augmented feedback for enhanced skill acquisition in individuals with traumatic brain injury. *Perceptual and Motor Skills.* 1996;82:507-514.
- 78. Wulf G, Shea CH. Understanding the role of augmented feedback: The good, the bad, and the ugly. In: Williams AM, Hodges NJ, eds. *Skill acquisition in sport: Research, theory and practice*. London: Routledge; 2004:121-144.

28

- 79. Wulf G, Shea CH. Principles derived from the study of simple motor skills do not generalize to complex skill learning. *Psychometric Bulletin and Review*.
 2002;9:185-211.
- Fasoli SE, Trombly CA, Ticle-Degned L, Verfaellie MH. Effect of instructions on functional reach in persons with and without cerebrovascular accident. *American Journal of Occupational Therapy*. 2002;56(4):380-390.
- 81. Landers M, Wulf G, Wallman H, Guadagnoli MA. An external focus of attention attenuates balance impairment in Parkinson's disease. *Physiotherapy*.in press.
- Totsika V, Wulf G. The influence of external and internal foci of attention in transfer to novel situations and skills. *Research Quarterly for Exercise and Sport*. 2003;74:220-225.