



NOVA

University of Newcastle Research Online

nova.newcastle.edu.au

Stevens, C. J., Bennett, K. J. M. & Novak, A. D. et al. (2019) Cycling power profile characteristics of national-level junior triathletes, *Journal of Strength and Conditioning Research*, 33(1) pp197-202

Available from: <http://dx.doi.org/10.1519/JSC.0000000000001876>

This is a non-final version of an article published in final form in the *Journal of Strength and Conditioning Research*, 33(1) pp197-202

Accessed from: <http://hdl.handle.net/1959.13/1416708>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Title: The cycling power profile characteristics of national level junior triathletes

Running head: Power profile characteristics of junior triathletes

Authors: Christopher J Stevens¹, Kyle J M Bennett², Andrew R Novak³, Aden B Kittel³ and Ben J Dascombe⁴

Affiliations

¹*School of Health and Human Sciences, Southern Cross University, Coffs Harbour, Australia*

²*Sport and Exercise Science, Faculty of Health, University of Technology Sydney, Moore Park, Australia*

³*Applied Sports Science and Exercise Testing Laboratory, Faculty of Science and Information Technology, University of Newcastle, Ourimbah, Australia*

⁴*Department of Rehabilitation, Nutrition and Sport, La Trobe University, Bundoora, Australia*

Address for correspondence

Dr Christopher John Stevens
School of Health and Human Sciences
Southern Cross University
Hogbin Dr
Coffs Harbour, 2450, NSW, Australia
Email: Christopher.Stevens@scu.edu.au
Ph: 0411 797 245

1 **ABSTRACT**

2 With the draft-legal rule recently introduced to junior triathlon competition, it has become
3 difficult to assess cycling performance through race results. Therefore, this study assessed the
4 cycling power profile characteristics of national level junior triathletes to assist with physical
5 assessment and program design. Thirteen male (17.0 ± 1.0 yr) and eleven female (17.2 ± 1.3
6 yr) national level junior triathletes completed a cycling power profile that consisted of
7 maximal intervals that lasted 6, 15, 30, 60, 240 and 600 seconds in duration. Each power
8 profile was completed on a LeMond ergometer using the subject's own bicycle, with power
9 output and cadence recorded for all intervals. Mean power output values for males ($783 \pm$
10 134 , 768 ± 118 , 609 ± 101 , 470 ± 65 , 323 ± 38 , 287 ± 34 W) were significantly ($P < 0.05$)
11 higher than females (554 ± 92 , 510 ± 89 , 437 ± 75 , 349 ± 56 , 248 ± 39 , 214 ± 37 W) across
12 all intervals, respectively. Peak power output values for males across the 6 and 15 second
13 intervals (1011 ± 178 and 962 ± 170 W) were also significantly higher than for females (674
14 ± 116 and 624 ± 114 W), respectively ($P < 0.05$). Developing junior triathletes should aim to
15 increase their capacity across the power profile above the mean values listed. Athletes should
16 further aim to have power outputs equal to that of the best performers and beyond to ensure
17 that they can meet the demands of any competition situation.

18
19 **Keywords:** triathlon, youth, draft-legal, coaching, testing

20
21 **INTRODUCTION**

22 Triathlon is a multidisciplinary sport encompassing the sequential completion of swimming,
23 cycling and running stages. In elite senior and junior competition, racing is classed as 'draft-
24 legal', permitting athletes to closely follow one another (i.e. drafting) during the cycling stage
25 to reduce drag forces (2, 11). While drafting may also be beneficial during the swimming and
26 running stages, it has particular importance during the cycling stage due to the increased wind

1 resistance creating greater drag at high speeds (12). Specifically, drafting behind small (i.e. 1-
2 4 riders) and large (i.e. 8 or more riders) groups of cyclists has been shown to reduce the
3 oxygen consumption requirement to sustain a given speed by as much as $27 \pm 7\%$ and $39 \pm$
4 6% , respectively (5). Hence, drafting allows individual competitors to alternate between
5 higher intensity efforts whilst leading the group or making a breakaway manoeuvre, with
6 interspersed lower intensity efforts when drafting to conserve energy. A study of male
7 international Olympic distance triathlon competition revealed that 34 ± 14 high intensity
8 efforts (>600 W) were performed during the cycling stage and 18% of total cycling time
9 exceeded maximal aerobic power (3), highlighting the intermittent demands of the race.
10 Hence, the tactical nature of drafting transforms the demands of the cycling stage into a high-
11 intensity, intermittent activity.

12
13 Due to the tactical nature of the draft-legal format, the cycling performance of opponent
14 triathletes during such competitions (i.e. their maximal performance over various durations)
15 is difficult to assess. Performance in the swimming and running stages can be inferred from
16 race times due to these stages more closely reflecting an individual time trial. However, in the
17 cycling stage, athletes take advantage of the draft effect and ride together in groups, which
18 means that they often finish with the same time (2). Also, many athletes will attempt to
19 minimise power output during the cycling stage in order to conserve energy prior to the
20 running stage (2, 8). Therefore, the optimal way to assess the maximal cycling capability of
21 an athlete over various durations is through controlled laboratory testing.

22
23 Current laboratory-based research on cycling in triathlon has focused on assessing maximal
24 aerobic capacity using incremental test protocols, with values as high as 74.3 ± 4.3 mL·kg⁻¹·
25 min⁻¹ reported for elite senior competitors (6). Further, maximal aerobic power values of

1 385-389 W have been reported for senior elite triathletes (4, 6). The application of such data
2 for draft-legal races is questionable considering that the high intensity, intermittent profile of
3 the draft-legal format requires the assessment of a triathlete's complete aerobic and anaerobic
4 capacities across various durations (3, 8). The cycling power profile is a reliable performance
5 test incorporating maximal self-paced intervals of 6-600 seconds in duration (8) and it has
6 recently been demonstrated to predict road cycling performance (8). It has also been
7 recommended by the Australian Institute of Sport as a useful cycling test protocol for
8 triathletes (13) and as a result it has been adopted by Australian state-level junior
9 representative triathlon squads. As such, this test has become important for physical
10 assessment and program design for these junior athletes, however, no normative data
11 currently exists for this population, which would likely assist coaches and athletes with their
12 interpretation of test results. Therefore the purpose of this study was to describe the
13 laboratory power profile results of junior male and female triathletes competing at the
14 national level.

15

16 **METHODS**

17 **Experimental Approach to the problem**

18 This descriptive study measured the power profile performance of national level junior
19 triathletes in a standardised laboratory test consisting of six maximal self-paced intervals (6,
20 15, 30, 60, 240 and 600 s in duration) with periods of active recovery (174, 225, 330, 480 and
21 600 s in duration) as described previously (8). All cycling was completed on each subjects's
22 own personal road bicycle that was attached to a LeMond Revolution cycle ergometer
23 (LeMond Fitness Inc., Woodinville, Washington, USA). The LeMond Revolution takes the
24 place of the rear wheel, using the bicycle's normal drivetrain to adjust resistance, which
25 allows the use of equipment and bicycle geometry that is specific to each individual. Power

1 output obtained from the LeMond Power Pilot (LeMond Fitness Inc., Woodinville,
2 Washington, USA) has previously been validated against the SRM power meter with the
3 level of agreement considered acceptable (7). Data was collected during training camps
4 leading into competition when the athletes were close to their peak condition.

5

6 **Subjects**

7 Thirteen male (age: 17.0 ± 1.0 yr, stature: 176.6 ± 5.7 cm, body mass: 65.8 ± 7.1 kg, sum of 7
8 skinfolds: 49.4 ± 10.2 mm, body fat: $8.7 \pm 1.7\%$) and eleven female (age: 17.2 ± 1.3 yr,
9 stature: 166.8 ± 7.9 cm, body mass: 57.5 ± 7.7 kg, sum of 7 skinfolds: 76.5 ± 15.5 mm, body
10 fat: $16.8 \pm 3.9\%$) national level junior triathletes volunteered for the study. Inclusion criteria
11 stipulated that subjects must be aged 16-19 years and currently competing in the Australian
12 National Junior Triathlon Series over the sprint distance (i.e. 750 m swim, 20 km cycle, 5 km
13 run). All subjects were familiar with riding on a cycle ergometer. All subjects and their
14 guardians provided written informed consent prior to testing. An institutional ethics
15 committee granted approval for the project (XXX H-2011-0350).

16

17 **Procedures**

18 An anthropometric profile was obtained from each participant consisting of stature (217
19 Stadiometer, Seca, Birmingham, United Kingdom), body mass (DS-530 electronic scales,
20 Wedderburn, Sydney, Australia) and skinfold thickness at seven sites (Harpenden Calipers,
21 Baty International, West Sussex, United Kingdom). The seven sites included bicep, tricep,
22 subscapular, supraspinalae, abdominal, quadriceps and medial calf and these sites were
23 summed to form the sum of 7 skinfolds (X_1). Body density was calculated with specific
24 regression equations for male (14) and female (15) Australian athletes as per below (where

1 X_2 = the sum of 6 skinfolds as above minus the bicep). Percent body fat was also estimated
2 via the equation below (9).

3

$$4 \quad \text{Male Body Density (14)} = 1.0988 - 0.0004(X_1)$$

$$5 \quad \text{Female Body Density (15)} = 1.20953 - 0.08294(\text{Log}_{10}X_2)$$

$$6 \quad \% \text{ Body Fat (9)} = [4.95/\text{Body Density} - 4.5] \times 100$$

7

8 For 24 hours prior to the power profile, caffeine and high intensity exercise were not
9 permitted and the athletes were instructed to consume their usual pre-race diet. The
10 participants performed a standardised 10 min warm-up that consisted of riding between 100-
11 200 W, as well as three six second intervals at 70, 80 and 90% of their perceived maximal
12 intensity, respectively. The power profile test commenced two minutes later and all intervals
13 began from a rolling start between 70-80 $\text{r}\cdot\text{min}^{-1}$. Verbal encouragement was provided during
14 the intervals and participants were instructed to self select and adjust their gear ratio at any
15 time to produce their best performance over each interval. The athletes were also instructed
16 that the shorter intervals (6-15 s) were a maximal sprint while the longer intervals (30-600
17 seconds) required a self-selected pacing strategy to produce the maximal mean power. During
18 active recovery, cyclists were instructed to pedal at a power output of <100 W. A 50
19 centimetre fan was placed 1 metre in front of the participant and provided a wind speed of 8
20 $\text{m}\cdot\text{s}^{-1}$ to simulate the convective cooling of outdoor conditions and tepid water (20-23°C) was
21 ingested *ad libitum* as recommended (10).

22

23

24

25

1 **Measures**

2 Power output and cadence were recorded at a frequency of 1 Hz using a LeMond Power
3 Pilot. The first second of data obtained in the 6 second intervals was not included in the data
4 analysis as per previous research (8). Heart rate was recorded with a Garmin Forerunner
5 910XT heart rate monitor wrist watch and chest strap (Garmin Ltd., Canton of Schaffhausen,
6 Switzerland). All data was downloaded post-test and arranged in Microsoft Excel (Microsoft
7 CorporationTM, Redmond, WA, USA) before further analysis. Power output data were also
8 divided by the participant's body mass to calculate relative values.

9

10 **Statistical Analyses**

11 The data were examined for assumptions of normality using the Kolmogorov-Smirnov test
12 and visually inspected through histograms and box plots. A two-way repeated measures
13 ANOVA was used to determine the main effects of sex on power output, cadence and heart
14 rate for each interval where it was measured. *Post hoc* comparisons with Bonferonni
15 adjustment were used to identify any significant differences. All statistical analysis were
16 conducted using SPSS software V22.0 (IBM Corporation, Somers, NY, USA). Power curves
17 were plotted for each athlete and group means using Microsoft Excel's built-in power
18 function ($R^2 > 0.94$ for all power curves) and a 'best performer' for both sexes was identified
19 as the athlete who achieved the highest power output across all interval durations in the
20 power profile itself and does not necessarily reflect the best performing triathlete in
21 competition.

22

23

24

25

1 **RESULTS**

2 The descriptive statistics for mean power output measures of the group and the best
3 performer across the power profile are presented in Table 1. All mean power outputs reported
4 were significantly higher in males than females for both absolute and relative measures
5 ($P<0.05$). Power curves of the group means and best performing male and female athlete
6 across the power profile tests are presented in Figure 1.

7
8 ***Insert Table 1 Here***

9 ***Insert Figure 1 Here***

10

11 The descriptive statistics for peak power output measures of the group and the best performer
12 across the 6 second and 15 second intervals are presented in Table 2. These peak power
13 outputs were both significantly higher in males when compared to females for both absolute
14 and relative measures ($P<0.05$).

15

16

Insert Table 2 Here

17

18 Mean and peak cadence measures of the group and best performer are presented in Table 3.
19 Mean cadence measures were significantly higher in males when compared to females across
20 the 15 and 30 second intervals ($P<0.05$). Peak cadence measures were significantly higher in
21 males when compared to females across the 6 and 15 second intervals ($P<0.05$). There were
22 no significant differences in cadences across any other interval ($P>0.05$).

23

24

Insert Table 3 Here

25

1 Mean heart rates across the 240 and 600 s intervals were 172 ± 8 beats·min⁻¹ and 179 ± 6
2 beats·min⁻¹ as well as 174 ± 7 beats·min⁻¹ and 178 ± 5 beats·min⁻¹ for males and females,
3 respectively. No significant differences were observed between sexes for the heart rate
4 measures ($P>0.05$).

5

6 **DISCUSSION**

7 This investigation has provided a novel insight into the cycling capacities of national level
8 junior triathletes. This information is useful for a number of purposes including the
9 preparation of athletes, monitoring changes in performance and talent identification. Such
10 data provides a set of normative values for regular cycle-based testing, which can also help to
11 identify specific strengths and weaknesses to benefit training prescription. Overall, the males
12 outperformed the females, even when corrected for differences in body mass, although the
13 gap between relative data for males and females was somewhat reduced. Further, males and
14 females employed significantly different cadences for the intervals shorter than 60 seconds
15 duration, however both cadences and physiological intensities were similar for the longer
16 duration intervals.

17

18 The power output requirements of the cycling stage within draft-legal junior triathlon are
19 highly variable, with the employed race tactics depending on a wide range of variables (2). In
20 addition, each course is highly variable, consisting of an entirely different circuit profile.
21 Therefore it is not adequate to prepare for such a race in this competition by simulating a
22 previous race in training (i.e. with the aid of performance times or race power outputs
23 through power meter analysis). Instead, developing junior triathletes should aim to be
24 physically superior by improving their capability to produce power across both aerobic and
25 anaerobic intervals (8), which is of high importance to draft-legal triathlon racing (3). The

1 current study described the mean cycling power outputs of junior triathletes in the power
2 profile, but also highlighted the power outputs of the best performer for both sexes.
3 Therefore, the current data should be used as a set of normative values for regular cycle-
4 based testing in these developing athletes. Developing junior triathletes and their coaches
5 should aim initially to have power outputs similar to the group mean. Secondly, athletes
6 should aim to produce power outputs equal to that of the best performer and beyond, which
7 would ensure that they can meet the demands of any competition situation and a greater
8 opportunity for successful performance.

9
10 The use of the power profile test combined with the the data in the current study may help an
11 athlete to identify specific weaknesses in their cycling ability. Such an example may be
12 where an athlete performs well relative to their peers in the longer intervals but does not
13 possess the anaerobic power to perform well in the short duration intervals. This result would
14 highlight the need for more maximal sprint training and perhaps resistance training exercises
15 which also serves to improve cycling sprint performance (16). Another advantage of regular
16 power profile testing is that the results can be useful for a coach to construct an informed
17 training program for an athlete in relation to their current level of fitness.

18
19 Along with a set of normative values for athletes and coaches to utilise, this study provides
20 normative cycling power functions (see equations in Figure 1) for high performing junior
21 triathletes. These power functions have a useful application for training and performance
22 testing and have not previously been reported for such a cohort. Importantly, the power
23 function allows for estimation of power outputs across any duration not explicitly assessed
24 within the test protocol or for individuals that have not undertaken a power profile. By simply
25 inserting the 'x' value of the duration of interest, the power functions provided can be used to

1 estimate normative maximal mean power output across any duration between 5–600 seconds.
2 Data may also be extrapolated beyond these limits if desired, for example, comparisons of
3 functional threshold power across 20 or 60 minutes (1) would require insertion of an ‘x’
4 value of 1200 or 3600, respectively. However, it should be noted that estimates may become
5 increasingly inaccurate for durations that lie further from the explicitly measured 5-600
6 second efforts of the power profile. Nevertheless, such estimates have strong implications for
7 coaches who may be limited for time within training camps and cannot conduct a power profile
8 assessment for 50 minutes with each individual athlete. Instead, the coach may choose
9 several efforts of any duration and compare these to the normative power functions ($W \cdot kg^{-1}$)
10 established in the current study. Coaches and athletes also have the option to compare
11 recordings from their mobile power meters during field-based training and/or during races,
12 with the normative power functions established in this study.

13
14 The power outputs were significantly higher in males compared to females and these
15 differences still existed after adjustments for body mass. Interestingly, mean and peak
16 cadences were significantly lower in females compared to males for most intervals lasting
17 less than 60 seconds. Considering gears were able to be freely selected by the athletes, this
18 suggests that the females preferred to perform shorter intervals at a lower cadence compared
19 to the males. It is difficult to speculate if the males would have performed better in a gear
20 with more resistance, or if the females would have performed better in a gear with less
21 resistance. In contrast, males and females chose a similar cadence in all of the intervals
22 lasting 60 seconds or longer. Also, mean heart rates were similar between the sexes across the
23 longer duration efforts, suggesting both sexes self-selected similar relative cycling intensities.

24

1 An important limitation of this study was that the study population consisted of only one fifth
2 of the triathletes competing in the Australian National Junior Triathlon Series. Indeed, a
3 larger sample size would make for a stronger set of normative data. Nevertheless, the current
4 study contained a broad spectrum of athletes, including the complete squad of two state
5 triathlon bodies. The study also includes both males and females who have gone on to
6 compete in the under 23 world triathlon championships and the senior elite category of the
7 International Triathlon Union World Triathlon Series. Hence, coaches can have confidence
8 that the data presented on the best performing athletes were of a high standard, however,
9 there may be better performing athletes who could not be included in this study. Another
10 limitation of the study was that the power profile protocol measured the power outputs from a
11 rested state, rather than a fatigued state, which would be more specific to a triathlon scenario.
12 The ability to perform anaerobic efforts under fatigue would be another useful indication of a
13 draft-legal triathlete's cycling ability.

14

15 **PRACTICAL APPLICATIONS**

16 The data described herein can be used as a set of normative values and normative power
17 functions for developing elite junior triathletes with the goal to perform well in draft-legal
18 competitions. With both the mean and best performing male and female power outputs and
19 resultant power functions clearly defined across the power profile, athletes can use these
20 values and/or equations as a training goal, or to help them identify their strengths and
21 weaknesses relative to their peers, which will be useful to inform training prescription.
22 Overall, it allows informed, evidence based decisions to be made by technical and
23 conditioning coaches in regard to the interpretation of cycling assessment and the cycling
24 program design of national level junior triathletes.

25

1 **Acknowledgements:** There was no outside financial support for this project. The authors
2 would like to acknowledge the subjects for their contribution to the study.

3

4 **Conflict of Interest:** There is no conflict of interest pertaining to the published data.

5

6 **REFERENCES**

- 7 1. Allen H and Coggan A. *Training and Racing with a Power Meter*. Boulder, CO:
8 VeloPress, 2006.
- 9 2. Bentley DJ, Millet GP, Vleck VE, and McNaughton LR. Specific aspects of
10 contemporary triathlon: Implications for physiological analysis and performance.
11 *Sports Med* 32: 345-359, 2002.
- 12 3. Etxebarria N, D'Auria S, Anson JM, Pyne DB, and Ferguson RA. Variability in power
13 output during cycling in international Olympic-distance triathlon. *Int J Sports Physiol*
14 *Perform* 9: 732-734, 2014.
- 15 4. Hue O. Prediction of drafted-triathlon race time from submaximal laboratory testing
16 in elite triathletes. *Can J Appl Physiol* 28: 547-560, 2003.
- 17 5. McCole SD, Claney K, Conte JC, Anderson R, and Hagberg JM. Energy expenditure
18 during bicycling. *J Appl Physiol* 68: 748-753, 1990.
- 19 6. Millet GP and Bentley DJ. The physiological responses to running after cycling in
20 elite junior and senior triathletes. *Int J Sports Med* 25: 191-197, 2004.
- 21 7. Novak AR, Stevens CJ, and Dascombe BJ. Agreement between LeMond Revolution
22 cycle ergometer and SRM power meter during power profile and ramp protocol
23 assessments. *JSC* 4: 37, 2015.
- 24 8. Quod MJ, Martin DT, Martin JC, and Laursen PB. The power profile predicts road
25 cycling MMP. *Int J Sports Med* 31: 397-401, 2010.
- 26 9. Siri WE. The gross composition of the body. *Adv Biol Med Phys* 4: 239-280, 1956.

- 1 10. Stevens C and Dascombe B. The reliability and validity of protocols for the
2 assessment of endurance sports performance: An updated review. *Meas Phys Educ*
3 *Exerc Sc* 19: 177-185, 2015.
- 4 11. Suriano R and Bishop D. Physiological attributes of triathletes. *J Sci Med Sport* 13:
5 340-347, 2010.
- 6 12. Swain DP. The influence of body mass in endurance bicycling. *Med Sci Sports Exerc*
7 26: 58-63, 1994.
- 8 13. Tanner R and Gore C (Australian Institute of Sport). *Physiological Tests for Elite*
9 *Athletes*. Lower Mitcham South Australia: Human Kinetics, 2013.
- 10 14. Withers RT, Craig NP, Bourdon PC, and Norton KI. Relative body fat and
11 anthropometric prediction of body density of male athletes. *Eur J Appl Physiol Occup*
12 *Physiol* 56: 191-200, 1987.
- 13 15. Withers RT, Norton KI, Craig NP, Hartland MC, and Venables W. The relative body
14 fat and anthropometric prediction of body density of South Australian females aged
15 17-35 years. *Eur J Appl Physiol Occup Physiol* 56: 181-190, 1987.
- 16 16. Yamamoto LM, Klau JF, Casa DJ, Kraemer WJ, Armstrong LE, and Maresh CM. The
17 effects of resistance training on road cycling performance among highly trained
18 cyclists: A systematic review. *J Strength Cond Res* 24: 560-566, 2010.

19
20 **Figure Captions**

21
22 **Figure 1:** Power curves and power functions of the group means and best performing male
23 and female athlete across the power profile tests.

24

Table 1. Mean power output measures of the group mean and the best performer expressed in both absolute and relative terms.

	Interval (s)	Group (W)	Best (W)	Group (W·kg ⁻¹)	Best (W·kg ⁻¹)
<i>Males</i>	6	783 ± 134	1000	11.9 ± 1.9	15.7
	15	768 ± 118	920	11.7 ± 1.4	14.5
	30	609 ± 101	761	9.2 ± 1.1	12.0
	60	470 ± 65	519	7.2 ± 0.8	8.2
	240	323 ± 38	333	4.9 ± 0.4	5.2
	600	287 ± 34	321	4.4 ± 0.4	5.0
<i>Females</i>	6	554 ± 92*	697	9.7 ± 1.2*	10.8
	15	510 ± 89*	654	8.9 ± 1.1*	10.1
	30	437 ± 75*	550	7.6 ± 0.9*	8.5
	60	349 ± 56*	455	6.1 ± 0.8*	7.0
	240	248 ± 39*	302	4.4 ± 0.7*	4.7
	600	214 ± 37*	271	3.8 ± 0.6*	4.2

Data is presented as mean ± standard deviation. s = seconds, W = watts, W·kg⁻¹ = watts per kilogram of body mass. *Significantly ($P < 0.05$) lower than males for respective interval duration.

Table 2. Peak power output measures of the group mean and best performer expressed in both absolute and relative terms.


	Interval (s)	Group (W)	Best (W)	Group (W·kg ⁻¹)	Best (W·kg ⁻¹)
<i>Males</i>	6	1011 ± 178	1346	15.3 ± 1.9	19.3
	15	962 ± 170	1234	14.6 ± 2.1	17.7
<i>Females</i>	6	674 ± 116*	864	11.8 ± 1.6*	13.4
	15	624 ± 114*	796	10.9 ± 1.4*	12.3

Data is presented as mean ± standard deviation. s = seconds, W = watts, W·kg⁻¹ = watts per kilogram of body mass. *Significantly ($P < 0.05$) lower than males for respective interval duration.

Table 3. Mean and peak cadence measures of the group mean and best performer.

	Interval (s)	Mean: Group (r·min⁻¹)	Mean: Best (r·min⁻¹)	Peak: Group (r·min⁻¹)	Peak: Best (r·min⁻¹)
<i>Males</i>	6	100 ± 9	92	118 ± 11	124
	15	112 ± 12	110	122 ± 17	161
	30	113 ± 10	113		
	60	109 ± 11	113		
	240	103 ± 11	113		
	600	98 ± 10	101		
<i>Females</i>	6	93 ± 11	98	108 ± 10*	116
	15	103 ± 8*	110	110 ± 10*	119
	30	102 ± 9*	108		
	60	103 ± 5	103		
	240	99 ± 7	100		
	600	99 ± 6	97		

Data is presented as mean ± standard deviation. r·min⁻¹ = revolutions per minute, s = seconds. *Significantly ($P < 0.05$) lower than males for respective interval duration. Peak cadence was not considered to be of relevance for intervals of >15 seconds.



ACCEPTED